



SOCIETY OF TRAUMA NURSES

THE ELECTRONIC LIBRARY OF

TRAUMA LECTURES

THE ELECTRONIC LIBRARY OF

TRAUMA LECTURES

The Trauma Team: From Prehospital Through the ED



SOCIETY OF TRAUMA NURSES

Objectives

At the conclusion of this presentation the participant will be able to:

- Describe the differences between Prehospital EMS and the Emergency Department (ED) environments
- Describe basic prehospital EMS operations
- Describe basic prehospital EMS care abilities
- Describe prehospital EMS to Trauma ED handoff
- Describe Trauma ED operations
- Recognize signs of trouble in the “fresh” trauma patient
- Describe evidence-based practice changes/controversies in prehospital and ED trauma care
- Explain tips to provide the best patient care

Underlying Principles

- The success of a trauma resuscitation is only as good as its team which includes providers from the prehospital environment through the ED Trauma Team
- Readiness, hyper-vigilance, consistent organization and clear-cut communication produces effective outcomes
- Practice makes perfect
- A proficient trauma team cannot function without a “skilled” team leader!

Side by Side Comparison


PREHOSPITAL EMS

- Protocol driven based on initial assessment
- Emphasis on rapid transport
- Limited resources

EMERGENCY DEPARTMENT

- Physician/Provider driven based on in-depth assessment
- Emphasis on treatment and resuscitation
- Vast resource availability

EMS Operations



EMS Operations


- Calls can be made any timeframe after a traumatic event occurs
- Dispatch centers are responsible for activating resources based on the information they receive
- Travel to the scene can take extended time



**911
Activation**

EMS Operations

- Preparations made based on information from dispatch center
- Air/MedEvac may be put on standby or launched
- Safe response is paramount
- Varying levels of care within the EMS system



Acuity of Patient	EMR	EMT/AEMT	Paramedic
Critical (Red)	Simple	Fundamental	Complex
Emergent (Yellow)		Simple	Fundamental
Low (Green)			Simple

Complex/Fundamental/Simple represent the depth of knowledge and skill each level of provider has for a particular acuity patient.

Levels of Care

EMR

- Simple lifesaving skills to provide immediate care to critical patients
- Limited equipment and education
- Cannot transport

EMT

- Basic, non-invasive, lifesaving care
- Basic equipment and basic education
- Can transport
- Comparable to an ED Tech

Levels of Care

AEMT

- Minimally invasive, low risk skills, lifesaving care
- Placement of airways
- IV/IO access with limited medication administration
- Enhanced equipment and advanced education
- Can transport

EMT-P

- Highest level of prehospital EMS care. Invasive skills with critical care level treatments
- High level equipment with extensive education building on the foundation of lower-level skills
- Can transport via ground and air

Levels of Care

Prehospital Hospital Providers

- RNs
- Advanced practitioners
- Physicians

(L-R) Dr. Stephan Russ, associate chief of staff for Vanderbilt University Medical Center and associate professor of Emergency Medicine, recently spent the day and flew with LifeFlight 3 (Clarksville) Nurses Jessica VanMeter, DNP, and Tony Smith, DNP



EMS Operations

- Scene size up
- Need for additional resources
- Primary Assessment of the patient
- Treatment of life-threatening injuries



**Initial
Assessment**

EMS Operations



- Symptoms
- Vital Signs
- SAMPLE Assessment
 - Symptoms
 - Allergies
 - Medications
 - Past Medical History
 - Last Oral Intake
 - Events

**Secondary
Assessment**

EMS Operations

- Trauma Criteria
- Spinal Stabilization
- Reassessment
- Destination & Notification
- On Scene Delays



A paramedic in a maroon shirt and blue gloves is kneeling and adjusting a white neck brace on a patient lying on a stretcher. The patient is wearing a plaid shirt and sunglasses. The stretcher is orange and has 'AMBULANCE' written on it. The scene is outdoors on a paved area with grass. A large orange bag is visible in the background.

Treatment

Delay at the Scene



Maintain C-spine/Open Airway



While sizing up the scene the prehospital provider determines if there is a reason to maintain C-spine on their patient

Immobilization Criteria

1. Altered Mental Status for any reason, including possible intoxication from alcohol or drugs (GCS < 15 or AVPU other than A)
2. Complaint of neck and/or spine pain or tenderness
3. Weakness, tingling, or numbness of the trunk or extremities at any time since the injury
4. Deformity of the spine not present prior to this incident
5. Distracting injury or circumstances (i.e. anything producing an unreliable physical exam or history)

High risk mechanisms of injury associated with unstable spinal injuries include, but are not limited to:

- Axial load (i.e. diving injury, spearing tackle)
- High speed motorized vehicle crashes or rollover
- Falls greater than standing height

Major Trauma Criteria

- GCS \leq 13
- Respiratory rate is < 10 or > 29
- Pulse rate < 50 or > 120
- Systolic BP < 90 mmHg
- Penetrating injuries to head, neck, torso or proximal extremities
- Two or more suspected proximal long bone fractures
- Suspected flail chest
- Suspected spinal cord injury or limb paralysis
- Amputation (except digits)
- Suspected pelvic fracture
- Open or depressed skull fracture
- Ejection/ partial ejection from vehicle
- Death in the same compartment
- Extrication time in excess of 20 minutes
- Vehicle collision resulting in 12 inches of intrusion in to the passenger compartment
- Motorcycle crash > 20 MPH or with separation of rider from motorcycle
- Falls from greater than 20 feet
- Vehicle rollover (90 degree vehicle rotation or more) with unrestrained passenger
- Vehicle vs. pedestrian or bicycle collision above 5 MPH

Pennsylvania Trauma Patient Destination Criteria

Physiologic Criteria:

- Patient does not follow commands (GCS Moto \leq 5)
- Hypotension, even a single episode (SBP < 90 mmHg)
- Respiratory rate < 10 or > 29 breaths/minute or need for ventilator support (< 20 in age < 1 year)

Anatomic Criteria:

- Penetrating injury to head, neck, torso, and extremities proximal to elbow or knee (unless obviously superficial)
- Chest wall instability or deformity (for example, flail chest)
- Two or more proximal lone-bone (humerus or femur) fractures
- Crushed/degloved/mangled or pulseless extremity
- Amputation proximal to wrist or ankle
- Pelvis fractures
- Paralysis (spinal cord injury)

YES

CATEGORY 1 TRAUMA

- Requires immediate transport to a trauma center (Level 1 or 2), if within 45 minutes
- Otherwise, transport to a Level 3 (preferred) or Level 4 trauma center if patient can arrive at the Level 3 or Level 4 center within 45 minutes or before an air ambulance can arrive to the patient's location
- Notify Trauma Center ASAP (including category and ETA)

Pennsylvania Trauma Patient Destination Criteria

Mechanism of Injury:

- Falls
 - Adult: > 20 feet (one story = 10 feet)
 - Children: > 10 feet or 2-3 x height of child
- High Risk Auto Crash
 - Passenger compartment intrusion, including roof: > 12 in. occupant site or > 18 in. into compartment at any site
 - Ejection (partial or complete) from automobile
 - Death in same passenger compartment
- Auto v. pedestrian/bicyclist thrown, run over, or significant (> 20 mph) impact
- Motorcycle crash > 20 mph

Other factors combined with traumatic injuries:

- Older Adults: SBP < 110 may indicate shock after age 65
- Anticoagulants or bleeding disorder
- Burns with trauma mechanism
- Pregnancy (> 20 weeks)
- Finger amputation

YES

CATEGORY 2 TRAUMA

EITHER:

- Contact Medical Command at closest Trauma Center (Level 1, 2, or 3) for authorization for air medical transport if needed.

OR

- Transport by ground to closest Trauma Center (Level 1, 2, or 3) (if within 45 minutes)
- Otherwise, transport to closest Level 4 Trauma Center (if within 45 minutes)

Pennsylvania Trauma Patient Destination Criteria

NO



CATEGORY 3 TRAUMA
TRANSPORT TO CLOSEST APPROPRIATE
RECEIVING FACILITY:

- Frequently reassess for Category 1 or 2 criteria
- Contact medical command, if doubt about appropriate destination

General Trauma Guidelines

Trauma Alert
Criteria

Scene Time

Destination
Protocols

Vascular
Access & Fluid
Administration

2011 CDC GUIDELINES

Physiologic
Criteria

Highest Level of Care
(Level 1 or 2 Trauma Center)

High Mechanism
of Injury
Indication

Highest Level of Care
(Level 1 or 2 Trauma Center)

2011 CDC Guidelines

Lower
Mechanism of
Injury

Closest Trauma Center
(regardless of level of care)

Special Patient
or System
Considerations

Possibly to a Trauma Center

EMS Operations

- Time to Transport:
 - 10 minutes or less if not entrapped
 - 20 minutes or less if entrapped
- Reasons to delay transport
 - Very few
- Enroute Treatments



**Rapid
Transport**

EMS Operations

- Arriving in the Trauma Bay
- Giving Report
- Assisting with Care
- Decon & Resupply



Arrival

Prehospital Care - Challenges



Environment
at the call



Prehospital Care Challenges

EMS services deemed essential in only 11 states

Unlike fire and police departments, EMS agencies are not considered an essential, or required, service in more than half of the country.



- Equipment
- Personnel experience
- Protocol limitations
- Unexpected events

**Prehospital Care -
Ventilation
*You Can (almost)
Always Bag***



Prehospital Care - Airway Adjuncts



King Airway



LMA



iGel



Combitube



Prehospital Care: Intubation

There have been numerous studies to evaluate the effectiveness of prehospital intubation.

Results range from improving to worsening outcomes for the patient.

Protocols vary state-by-state on how intubation can be performed.

EMS Intubation Protocol Sample



- Consider intubation if GCS is less than 8 or airway cannot be maintained.
- If patient is intubated or has an airway such as Combitube, King, LMA $P_{ET}CO_2$ levels should be continually monitored and maintained at 33 – 43 mmHg if available.

Prehospital Care – Hemorrhage Control

Proven effective techniques of hemorrhage control:

- Direct pressure
- Tourniquets

Unproven techniques:

- Elevation of limb
- Pressure points
- Cold application



Prehospital Care - Tourniquets

Special Operations Forces Tactical Tourniquet



Indications for Tourniquet


- Hemorrhage from an extremity that cannot be controlled with direct pressure or a pressure bandage
- Traumatic amputation
- There may be times (tactical, rescue, entrapped patient, multiple patient scene, backwoods environment) when tourniquet application is the best first option

There are no contraindications for applying a tourniquet

Tourniquet Safety



2125



Prehospital Care – Intravascular Access

2 large bore IV's?

Common EMS Protocol

- *Do not delay transport to establish IV/IO access.*
- Initiate IV/IO (18ga or larger) and hang NS, if approved.
- Consider 2nd IV/IO where hypovolemia is suspected (Adult only)
- (Adult) If SBP < 100 mmHg or heart rate > 120, initiate a fluid bolus of Normal Saline: 500 ml

Prehospital Care – Intravascular Access via Intraosseous Access



VIEWER DISCRETION IS ADVISED

<https://youtu.be/SYfVFq0rc7c>

EZ-IO[®]
Intraosseous Infusion System



Prehospital Care - Splinting



Prehospital Care – Splinting

Traction Splints



Prehospital Care – Splinting


Pelvic Fractures

- Physical exam is unreliable
- DO NOT ROCK or palpate the pelvis in the prehospital arena
- Avoid log rolling as much as possible
- Apply splint if in your area protocols



www.expeditionmedicine.co.uk/

Trauma ED Operations



Trauma ED Operations

- Notification received from EMS
- Information interpreted and plan is developed
- Questions asked

**EMS
Notification**



Yesterday 12:16 AM

Trauma ALERT FALL ETA: 10 mins

Trauma Alert GSW ETA: now

Yesterday 3:00 AM

Trauma Alert GSW ETA: now

Trauma Response GSW ETA: 5mins

Yesterday 6:21 AM

Trauma ALERT FALL ETA: now

Trauma Response FALL ETA: now

Trauma ED Operations

- Method of Trauma Alert Activation
- Personnel at Trauma Alert Activation
- Clearing Trauma Alert Activation

**Trauma
Activation**

Trauma ED Operations

- First ED encounter with the patient
- Report is given and questions are asked
- Care of the patient begins



**EMS Arrival
& Report**

ED Patient Care

- A-B-Cs
- Primary Assessment
- Imaging / Procedures
- Reassessment



We All Need to Know Our A, B, C, Ds

1. All trauma patients need oxygen until proven otherwise
2. All trauma patients are bleeding until proven otherwise
3. All trauma patients have a cervical spine injury until proven otherwise
4. All unconscious trauma patients have a brain injury until proven otherwise



Airway

Hemodynamically unstable trauma patients need an early definitive airway

Prevents aspiration

Reduces the oxygen debt of breathing

Maximizes systemic oxygen delivery


Corrects acid base disturbances

Allow clinician to focus on other life-threatening issues



BREATHING

All trauma patients need oxygen until proven otherwise



A photograph of a trauma scene. In the foreground, there is a large, dark red pool of blood on a light-colored floor. To the left, the lower legs and feet of several people wearing green protective suits and boots are visible. In the center and right, there is medical equipment, including a white corrugated tube and a blue bag. The background is slightly out of focus, showing more of the scene.

CIRCULATION / HEMORRHAGE


All trauma
patients are
bleeding
until
proven
otherwise

Hemorrhage: Five Anatomic Areas of Origin

Anatomic site	Modality	Reliability
Intraperitoneal	FAST	Positive test is diagnostic; negative test is suspect
	DPL	Gross positive is diagnostic; technically positive only is suspect
Retroperitoneal	Pelvic radiograph	Unstable fracture pattern is suggestive
Thoracic (pleural)	Chest radiograph	Positive test is usually diagnostic
Multiple long-bone fractures	Physical examination	US and radiograph confirmatory but do not quantify blood loss
External bleeding	Physical examination	Positive findings diagnostic but do not quantify blood loss

Neurological

All
unconscious
trauma patients
have brain injury
until proven
otherwise.



Most Potentially Preventable Trauma Deaths Are Related to:

- Airway obstruction
- Hemorrhage
 - Hemopneumothorax
 - Intracavitary bleeding
 - Intracranial hemorrhage



Situation Awareness


➤ Cause of instability must be recognized and corrected quickly by using a systematic approach.

➤ It is important to identify and prioritize systemic compromise.

**Detection
(Perception)**

Diagnosis

Prediction




Recognize Trouble!

Episodic Hypotension

Trauma patients with prehospital hypotension that are normotensive on arrival should have an arterial blood gas with Base Deficit (BD) interpreted early upon admission to help identify those patients who are at risk for “crumping.”


Routine Chest X-ray



Chest x-ray

- Early
- Valuable
- Significant injury
- Position of patient
- Never delay procedures for x-ray

CT Scan for Thoracic Trauma



An occult pneumothorax seen on a CT scan that was not detected on a plain anteroposterior supine chest radiograph.

- Recognize the insensitivity of the AP CXR to detection of pneumothoraces
- Remarkably large occult pneumothoraces may be present without an obvious anterolateral pleural stripe on AP CXR

Routine Pelvic X-ray



ATLS advocates that a routine pelvic x-ray be an adjunct to primary survey in trauma patients with **multiple injuries.**


Routine Cervical Spine X-ray

- The primary screening modality is axial computed tomography (CT) from the occiput to T1 with sagittal and coronal reconstructions.
- Plain radiographs contribute no additional information and should not be obtained.



Computed Tomography

- Used for hemodynamically stable patients
- Noninvasive procedure



2017051120
10:18:45
Kem: B
CT
VR


Flow:
Col:
Stat

L

120.000 kV
311 mA
Tilt: 0.000



30.00 mm/k



ED Patient Care

- Resuscitation
- Bedside Invasive Procedures



Treatment

Resuscitation

- For Resuscitation to occur:
 - Adequate perfusion and tissue oxygenation must be restored
- Resuscitation measures directed at:
 - Control the bleeding
 - Maintaining or restoring
 - Hematocrit
 - Normothermia
 - Normal INR
 - Base deficit within normal limits



Recognize Trouble!



Indicators of Anaerobic Metabolism and Acidosis

↓ serum pH (7.35 – 7.45)

↑ base deficit (2 to -2)

↑ serum lactate (0.4 – 1.8 mmol/L)

lactic acidosis:



Increased Unmeasured Anions

Recognize Trouble!

Base Deficit

- Amount of fixed base (bicarbonate) required to correct the pH of an aliquot of blood to 7.40 at 37 C
- Excellent correlation of base deficit to lactate and pH
- Normalizes rapidly with adequate resuscitation and control of hemorrhage

Optimizing Resuscitation

Prevent Unnecessary Blood Loss

- Low volume blood sampling
 - 3 cc for ISTAT: Na/K/Cl/Ca, Hgb, Hct, PO₂, PCO₂, pH, PT, INR, Lactic acid, Creatinine, Troponin
- Use FAST to rapidly detect intra-abdominal fluid
- Auto transfusion of thoracic blood using a chest tube drainage/auto-transfusion system
 - Avoid spilling blood on floor during chest tube insertion



Optimizing Resuscitation

Controlling Hemorrhage

- Direct pressure to all external bleeding
- Rapid suturing of all scalp and facial wounds
- Rapid application or evaluation of traction splints for femur fracture
- Binder for pelvic fracture
- Rapid reduction and pressure dressings of mangled extremities
- Leaving prehospital tourniquets in place until ready for surgical intervention

Endpoints of Resuscitation



- B/P & HR WNL
- Increased urine output
- Improved base deficit
- Decreased serum lactate
- Improved serum pH
- Normothermia
- Adequate coagulation status

Damage Control Resuscitation

Lifesaving interventions aimed to prevent and/or rapidly treat blood loss, irreversible hypothermia, acidosis, and coagulopathy in the unstable trauma patient with severe injuries.

A = acidosis

B = blood loss

C = cold


D = damage

Interventions to Decrease the Need for Blood Transfusion

- Intervene early in resuscitation to STOP bleeding
- Conserve blood early
- Aggressive, rapid, and efficient operative interventions
- Use novel surgical methods to STOP bleeding in the operating theater
- Early use of interventional radiology when appropriate
- Correct hypothermia, acidosis, coagulopathy

Trauma ED Operations

Disposition




- Operating Room
- Interventional Radiology
- Intensive Care Unit
- Termination of Efforts

Disposition - OR

Operating Room

- Rapid exposure
- Evacuation of blood
- Packing
- Control contamination
- Anticipate and fight the “trauma triad” of death: Hypothermia, Acidosis, Coagulopathy

Reassess/Plan



Disposition - ICU





Recognize Trouble!

Persistent Metabolic Acidosis

Continued
bleeding


Inadequate
resuscitation

Myocardial
dysfunction

General Hospital

← **Emergency
& Shock
Trauma
Only**

**Providing the
Best Care
Leadership &
Teamwork**





Providing the Best Care Communication

Physiologic Criteria

Mechanism of Injury

Interventions

Providing the Best Care Trauma Team Training Course

Prepares teams of five to care for trauma victims with the limited resources found at African rural hospital and health centers



Safer Surgery, Obstetrics + **Communities in Africa**
Canadian Network for International Surgery

Providing the Best Care

Goal Directed Therapy

- High risk patients
 - Geriatric patients
 - Pediatric patients
 - Pregnant women
- Anticoagulants
- Beta blockers



Providing the Best Care

“Routine Trauma Labs”

Information provided by routine admission chemistry and coagulation profiles in trauma patients seldom lead to clinical interventions.

Should not be ordered routinely on admission in trauma patients



Providing the Best Care

EAST Practice Guidelines

“Withholding and termination of resuscitation of adult cardiopulmonary arrest secondary to trauma: Resource document to the joint NAEMSP-ASCOT position statements”



The Eastern Association for the Surgery of Trauma

Consistent Organization and Clear-cut Communication Produce Effective Outcomes



Well established and maintained clear roles and goals

Similar training- street through hospital

Effective function of the team begins with the Team Leader

Summary and Conclusions

- It takes a team.
- Efficient teams communicate and have a defined leader.
- Effective teams use research-based protocols.
- Trauma centers save lives!

The Trauma Team: From Prehospital Through the Emergency Department

1. As the prehospital provider approaches the scene of a trauma call, they perform:
 - a. A radio transmission to the hospital
 - b. A scene size up
 - c. An estimate of neck size for c-collar
 - d. An estimate of victim's height and weight

2. Field intubation has been proven to improve outcome in:
 - a. Patients with BP less than 90 mm Hg
 - b. Patients with GCS less than 9
 - c. Patients with acute respiratory distress
 - d. None of the above

3. A proven technique of hemorrhage control is:
 - a. Direct pressure
 - b. Elevate above the heart
 - c. Pressure points
 - d. Cold application

4. Prehospital care for apparent pelvic fractures includes:
 - a. DO NOT ROCK or palpate the pelvis in the prehospital arena
 - b. Avoid log rolling as much as possible
 - c. Apply splint if in your area protocols
 - d. All of the above

5. Most preventable deaths in trauma care are due to:
 - a. Delay in CPR
 - b. Cardiac tamponade
 - c. Airway obstruction
 - d. Tension pneumothorax

6. For resuscitation to occur, there must be:
 - a. Cellular perfusion and tissue oxygenation
 - b. Restoration of a blood pressure greater than 90mm Hg
 - c. A hemoglobin greater than 9g/dL
 - d. A PaO₂ greater than 80 mm Hg

7. The Trauma Triad of Death is:
 - a. Hypotension, tachycardia and decreased urine output
 - b. Infection, inadequate nutrition, DVT's
 - c. Hypothermia, acidosis, and coagulopathy
 - d. Splenic, pelvic, and head injuries

8. A plain pelvic x-ray should be performed on all:
 - a. Trauma patients
 - b. Patients with head injury
 - c. Patients with multiple injuries
 - d. Patients over 65 years old

9. Primary screening for cervical spine injuries is:
 - a. CT scan
 - b. Cross table lateral C-spine
 - c. Swimmer's view C-spine
 - d. MRI

10. Persistent metabolic acidosis in a trauma patient may indicate:
 - a. Hyperventilation
 - b. Continued bleeding
 - c. Severe head injury
 - d. Hyperthermia

The Trauma Team: From Prehospital Through the Emergency Department

1. As the prehospital provider approaches the scene of a trauma call, they perform:
 - a. A radio transmission to the hospital
 - b. A scene size up**
 - c. An estimate of neck size for c-collar
 - d. An estimate of victim's height and weight

2. Field intubation has been proven to improve outcome in:
 - a. Patients with BP less than 90 mm Hg
 - b. Patients with GCS less than 9
 - c. Patients with acute respiratory distress
 - d. None of the above**

3. A proven technique of hemorrhage control is:
 - a. Direct pressure**
 - b. Elevate above the heart
 - c. Pressure points
 - d. Cold application

4. Prehospital care for apparent pelvic fractures includes:
 - a. DO NOT ROCK or palpate the pelvis in the prehospital arena
 - b. Avoid log rolling as much as possible
 - c. Apply splint if in your area protocols
 - d. All of the above**

5. Most preventable deaths in trauma care are due to:
 - a. Delay in CPR
 - b. Cardiac tamponade
 - c. Airway obstruction**
 - d. Tension pneumothorax

6. For resuscitation to occur, there must be:
 - a. Cellular perfusion and tissue oxygenation**
 - b. Restoration of a blood pressure greater than 90mm Hg
 - c. A hemoglobin greater than 9g/dL
 - d. A PaO₂ greater than 80 mm Hg

7. The Trauma Triad of Death is:
- a. Hypotension, tachycardia and decreased urine output
 - b. Infection, inadequate nutrition, DVT's
 - c. Hypothermia, acidosis, and coagulopathy**
 - d. Splenic, pelvic, and head injuries
8. A plain pelvic x-ray should be performed on all:
- a. Trauma patients
 - b. Patients with head injury
 - c. Patients with multiple injuries**
 - d. Patients over 65 years old
9. Primary screening for cervical spine injuries is:
- a. CT scan**
 - b. Cross table lateral C-spine
 - c. Swimmer's view C-spine
 - d. MRI
10. Persistent metabolic acidosis in a trauma patient may indicate:
- a. Hyperventilation
 - b. Continued bleeding**
 - c. Severe head injury
 - d. Hyperthermia

References

The Trauma Team: From Prehospital Through the Emergency Department

5th edition

American College of Surgeons Committee on Trauma (2018). Advanced Trauma Life Support, ATLS Student Course Manual, 10th ed. Chicago, American College of Surgeons.

American College of Surgeons, Committee on Trauma (2014) Resources for Optimal Care of the Injured Patient, 6th ed. Chicago, American College of Surgeons.

Cotton, B. A., Jerome, R., Collier, B. R., Khetarpal, S., Holevar, M., Tucker, B., Kurek, S., Mowery, N. T., Shah, K., Bromberg, W., Gunter, O. L., Riordan, W. P., Jr, & Eastern Association for the Surgery of Trauma Practice Parameter Workgroup for Prehospital Fluid Resuscitation (2009). Guidelines for prehospital fluid resuscitation in the injured patient. *The Journal of trauma*, 67(2), 389–402. <https://doi.org/10.1097/TA.0b013e3181a8b26f>

Gnugnoli DM, Singh A, Shafer K. EMS Field Intubation. [Updated 2021 Dec 9]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2022 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK538221/>

Guidelines for Field Triage of Injured Patients (2011), published by the Centers for Disease Control and Prevention and retrieved from <https://www.cdc.gov/mmwr/preview/mmwrhtml/rr6101a1.htm>.

Leidel, B. A., Kirchhoff, C., Bogner, V., Stegmaier, J., Mutschler, W., Kanz, K. G., & Braunstein, V. (2009). Is the intraosseous access route fast and efficacious compared to conventional central venous catheterization in adult patients under resuscitation in the emergency department? A prospective observational pilot study. *Patient safety in surgery*, 3(1), 24. <https://doi.org/10.1186/1754-9493-3-24>

Loyd JW, Larsen T, Swanson D. Aeromedical Transport. [Updated 2021 Aug 11]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2022 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK518986/>

“STOP THE BLEED” campaign retrieved from <https://www.stopthebleed.org/>